

**NIGHTTIME COLD AND FLU MULTI SYMPTOM- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled**  
**Strategic Sourcing Services LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Safeway Nighttime Cold & Flu**

**Active Ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan Hydrobromide 15 mg

Doxylamine Succinate 6.25 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Antihistamine

**Uses**

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do

not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

### **Other Information**

- store between 15-30°C (59-86°F)
- avoid excessive heat

### **Inactive Ingredients**

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

\*may contain this ingredient

### **Questions or Comments?**

Call **1-877-753-3929 Monday-Friday 9AM-5PM EST**

### **Principal Display Panel**

Compare to Vicks® NyQuil® LiquiCaps® active ingredients†

Multi-symptom cold & flu

#### **night time liquid caps**

**Acetaminophen**, Dextromethorphan HBr, Doxylamine Succinate

Pain Reliever / Fever Reducer

Cough Suppressant / Antihistamine

SOFTGELS

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**Another Quality Product Distributed by McKesson**

**One Post Street, San Francisco, CA 94104**

**[www.healthmart.com/healthmartbrand](http://www.healthmart.com/healthmartbrand)**

**Package Label**

**McKesson**

This product is not manufactured or distributed by The Procter & Gamble Company, Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

**16 softgels** (\*\*Liqui-filled capsules)

actual size



pain reliever/fever reducer  
cough suppressant/antihistamine

# multi-symptom cold & flu night time liquid caps

acetaminophen, dextromethorphan HBr,  
doxylamine succinate

compare to Vicks® NyQuil®  
LiquiCaps® active ingredients†

**Health Mart** PHARMACY



NDC 62011-0262-1

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.  
TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

Lot No.:  
Exp. Date:

**Drug Facts****Active ingredients (in each softgel)****Purpose**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 15 mg.....Cough suppressant  
Doxylamine succinate 6.25 mg.....Antihistamine

**Drug Facts (continued)****Uses**

- temporarily relieves common cold and flu symptoms
- sore throat ■ headache ■ minor aches and pains
- fever ■ runny nose and sneezing
- cough due to minor throat and bronchial irritation

**Drug Facts (continued)****Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have: ■ liver disease ■ alcoholism

**Drug Facts (continued)****Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each

<p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>■ liver disease</li> <li>■ glaucoma</li> <li>■ trouble urinating due to an enlarged prostate gland</li> <li>■ a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema</li> <li>■ cough that occurs with too much phlegm (mucus)</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>■ taking the blood thinning drug warfarin</li> <li>■ taking sedatives or tranquilizers</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>■ do not exceed recommended dosage</li> <li>■ excitability may occur, especially in children</li> <li>■ avoid alcoholic drinks</li> <li>■ marked drowsiness may occur</li> <li>■ alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>■ be careful when driving a motor vehicle or operating machinery</li> </ul>	<p><b>label to ensure correct dosing</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 15-30°C (59-86°F)</li> <li>■ avoid excessive heat</li> </ul> <p><b>Inactive ingredients</b> D&amp;C yellow #10, edible white ink, FD&amp;C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan*, sorbitol</p> <p>*may contain this ingredient</p> <p><b>Questions or comments?</b> Call 1-877-753-3935 Monday-Friday 9AM-5PM EST</p>
--	--

## HealthMart NightTime Cold & Flu multi-symptom

## NIGHTTIME COLD AND FLU MULTI SYMPTOM

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:62011-0262
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	P30;94A;SCU1;215;P120
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0262-1	16 in 1 CARTON	12/31/2015	02/23/2024
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/31/2015	02/23/2024

**Labeler** - Strategic Sourcing Services LLC (116956644)

Revised: 11/2021

Strategic Sourcing Services LLC